



JUST PUBLISHED

Bryozoan Studies 2001 - Proceedings of the 12th International Bryozoology Association Conference, Dublin, Ireland, 16-21 July 2001

Edited by P.N. Wyse Jackson, C.J. Buttlar & M.E. Spencer Jones

A.A. Balkema, Lisse; ISBN: 90 5809 388 3; xii+409 pp.

PRICE: ₤ 99,00 (£57.69 (approx.); \$94.05 (approx)).

Please order copies direct from the publishers via their web page: <http://balkema.ima.nl>

Contents

Foreword

Dedication

- Bryozoan communities in the Weddell Sea, Antarctica: a first overview - *B. Bader*
- Modelling multivariate determinants of growth in Antarctic bryozoans - *D.K.A. Barnes & S. De Grave*
- Seasonality and inter-annual variability in recruitment patterns of temperate encrusting fauna - *D.K.A. Barnes & B.C. Maughan*
- Use of radioactive labelled food to assess the role of the funicular system in the transport of metabolites in the cheilostome bryozoan *Membranipora membranacea* (L.) - *M.A. Best & J.P. Thorpe*
- Use of radioactive labelled silt to show depletion of food supply by upstream colonies of *Flustrellidra hispida* (Fabricius) - *M.A. Best & J.P. Thorpe*
- Colony life then and now: Lower Paleozoic trepostomes (500-350 mya) and living cyclostomes: a review - *R.S. Boardman*
- Bryozoans from temperate Pleistocene deep-water mounds, Great Australian Bight, Australia - *Y. Bone & N.P. James*
- Observations on ecological adaptations of *Lanceopora smeatoni* (MacGillivray), West Island, South Australia - *K.M. Brown, R. Schmidt & Y. Bone*
- Larval release pattern in Antarctic bryozoans - *J.M. Cancino, F. Torres & H.I. Moyano G.*
- Asexual propagation in the cheilostomes *Metrarabdotos* and *Coscinopleura*: effects on genetic variation and larval productivity - *A.H. Cheetham*
- Origin and early development of the International Bryozoology Association - *A.H. Cheetham*
- Influence of colony morphology on associated biota diversity in four Bryozoa - *S. Cocito, F. Ferdeghini, S. Pisaroni & D. Bedulli*
- Identification key for North American Ordovician trepostome families - *R.J. Cuffey*
- Deconstructing bryozoans: origin and consequences of a unique body plan - *R.A. Dewel, J.E. Winston & F.K. McKinney*
- Colony morphologies and missed opportunities during the Cincinnati (Late Ordovician) bryozoan radiation: examples from *Heterotrypa frondosa* and *Monticulipora mammulata* - *J.M. Erickson & D.A. Waugh*
- Systematics and biogeography of the Permian bryozoans in Europe - *A. Ernst*
- Late Cretaceous-Paleocene "porinids" - mixed frontal shields and evidence of polyphyly - *D.P. Gordon*
- Kubaninella*: a new genus of Adeonidae (Bryozoa: Cheilostomata) from the Western Kamchatka shelf of the Sea of Okhotsk - *A.V. Grischenko & S.F. Mawatari*

- Partitioning phenotypic variation: implications for morphometric analyses (Bryozoa) - *S.J. Hageman, M. Bayer & C.D. Todd*
- Bryozoan distribution in a Messinian coral reef complex of western Algeria - *A. Hamdane & P. Moissette*
- Effects of sewage on bryozoan diversity in Mediterranean rocky bottoms - *J.G. Harmelin & S. Capo*
- Predominance of obligate outbreeding in the simultaneous hermaphrodite *Celleporella hyalina sensu lato* - *R.N. Hughes, P.J. Wright, P.H. Manríquez & J.D.D. Bishop*
- Functional morphology of maculae in a giant ramose bryozoan from the Permian of Greenland - *M.M. Key, Jr., L. Thrane & J.A. Collins*
- Freshwater bryozoan remains from the Molteno Formation (Upper Triassic) of South Africa - *R. Kohring & A. Hörnig*
- Bryozoa of the high Arctic fjord – a preliminary study - *P. Kuklinski*
- Chesterian (Carboniferous) *Septopora* (Order Fenestrata), eastern North America - *F.K. McKinney*
- The ctenostome collar – an enigmatic structure - *M.J. McKinney & R.A. Dewel*
- Mitochondrial evidence of geographical isolation within *Bugula dentata* Lamouroux - *J.A. Mackie, M.J. Keough, J.A. Norman & L. Christidis*
- Results of a freshwater bryozoan survey in the Pacific Northwestern United States - *T. Marsh & T.S. Wood*
- Highly diverse bryozoan faunas from the Plio-Pleistocene of the Greek island of Rhodes - *P. Moissette, N. Spjeldnaes & E. Georgiades-Dikeoulia*
- Interactions of Bryozoans and Microbes in a Chemosynthetic Hydrothermal Vent System: Big Cove Formation (Lower Codroy Group, Lower Carboniferous, Middle Viséan/Arundian), Port au Port Peninsula, Western Newfoundland, Canada - *P.A. Morris, P.H. von Bitter, P.E. Schenk & S.J. Wentworth*
- Bryozoa from Oceanic south eastern Pacific Islands: diversity and zoogeography - *H.I. Moyano G.*
- A Moscovian (Carboniferous) bryozoan buildup from Svalbard - *H.A. Nakrem*
- Brood chambers in cribrimorphs evolved by fusion of costae: further arguments- *A.N. Ostrovsky*
- Neurotoxin found in the freshwater bryozoan *Lophopodella carteri* - *N.L. Peterson*
- Bryozoans in Ordovician depositional sequences, Cincinnati Arch region, USA - *J.R.P. Ross & C.A. Ross*
- Bryobaculum carinatum* gen. et sp. nov.: a new Mediterranean Pliocene deep-water bryozoan - *A. Rosso*
- Life history characters and ecology of some encrusting ctenostomes - *J.S. Ryland*
- Eocene Bryozoan Assemblages of the St Vincent Basin, South Australia - *R. Schmidt & Y. Bone*
- Changing concepts in species diversity in the northeastern Pacific - *D.F. Soule, J.D. Soule & P.A. Morris*
- Brooding in the Cretaceous bryozoan *Stichomicropora* and the origin of ovicells in cheilostomes - *P.D. Taylor & F.K. McKinney*
- The effects of increased external Ca⁺⁺ and K⁺ concentrations on the waveform dynamics of bryozoan spermatozeugmata - *M.H. Temkin*
- New bryozoan faunas from the Miocene of Burgenland (Austria) - *N.R. Vávra*
- Functional morphology of the anastomosing frondose growth form reported in *Heterotrypa frondosa* (d'Orbigny) (Bryozoa: Trepostomata) from the Cincinnati (Late Ordovician) of Ohio - *D.A. Waugh & J.M. Erickson*
- Freshwater bryozoans: a zoogeographical reassessment - *T.S. Wood*
- The reproductive cycle of *Plumatella casmiana* (Phylactolaemata: Plumetellidae) - *E.R. Wöss*
- A comparison of the early astogeny and life history of *Celleporella carolinensis* and *Celleporella hyalina* - *P.J. Wright & R.N. Hughes*
- Palaeoenvironmental interpretation of the Tramore Limestone Formation (Llandeilo, Ordovician) based on bryozoan colony form - *P.N. Wyse Jackson, C.J. Butcher & M.M. Key, Jr.*
- An Upper Eocene bryozoan fauna from Perwang-1 borehole, Salzburg, Austria - *K. Zágorská*
- Author index to the conference volumes (1-12) of the International Bryozoology Association (1969-2002) - *P.N. Wyse Jackson*

List of participants

Taxonomic index