



Paper in:

Patrick N. Wyse Jackson & Mary E. Spencer Jones (eds) (2002) *Annals of Bryozoology: aspects of the history of research on bryozoans*. International Bryozoology Association, Dublin, pp. viii+381.

Historical review on individual autozooidal behaviour and feeding mechanisms in Bryozoa

Andrew N. Ostrovsky*, Natalia N. Shunatova and Irina I. Antipenko

*Department of Invertebrate Zoology, Faculty of Biology and Soil Science,
St. Petersburg State University, Universitetskaja nab. 7/9, St. Petersburg, 199034,
Russia*

**Present address: Institut für Geowissenschaften, Christian-Albrechts-Universität zu
Kiel, Olshausenstr. 40, 24118 Kiel, Germany*

1. Introduction
2. Eighteenth century studies
3. Nineteenth century studies
4. Twentieth century studies
5. Conclusions
6. Acknowledgements

1. Introduction

During our studies on bryozoan autozooidal behaviour^{1,2} we, following Winston, realized that ‘the 19th [as well as 18th] century knowledge of individual (polypide) behaviour was never codified’.³ This negligence in respect to the work of the old authors led to two main consequences. Firstly, the names of many brilliant naturalists, whose observations and detailed descriptions were, as a rule, perfect and whose results are the basement of modern knowledge, were either undeservedly forgotten or mentioned only in passing. Secondly, ‘double work’ was carried out since many of previously recorded phenomena and structures were newly redescribed, which is intimately connected with the problem of scientific priority. So, this paper was written with the aim to correct the situation in question at least partially. We present a historical survey on the development of views on individual autozooidal behaviour and feeding mechanisms in Bryozoa, with the emphasis on the priority in the recording of one or other type of activities and feeding/sensory structures.

2. Eighteenth century studies

The first description of bryozoan behaviour was given by Abraham Trembley,⁴ who

observed the phylactolaemate bryozoan *Lophophus*⁵ in 1741. A brief summary of Trembley's notes was published first by Henry Baker,⁶ who later made and published his own observations of phylactolaemates.⁷ They both observed polypide protrusion and retraction, water currents carrying small animals into the lophophore, various movements of the lophophore and tentacles connected with both feeding and particle rejection (for instance, Trembley recorded tentacle flicks, lophophore expansion, accompanying the particle rejection, and particle transference along the oral grooves of lophophore), and defecation. Trembley⁸ and later Baker attributed movements of water within and around the lophophore only to the perpetual swinging or 'vibrating motion' of the tentacles.⁹ In contrast to Trembley, who believed that the tentacles did not touch the particles and bent inside the lophophore in order to create turbulent currents, Baker described particle capture by tentacles when these particles had escaped from the feeding current. Judging from the description, he was the first who recorded this kind of feeding by tentacular flicks when the previous mechanical contact between tentacle and particle was absent.¹⁰ August Rösel von Rosenhof also worked on phylactolaemates and correctly remarked that during production of currents the tentacles were often motionless.¹¹ He supposed that the currents were produced by a stream issuing from the contracting mouth.

John Ellis was apparently the first to publish descriptions and drawings of living marine bryozoans, briefly mentioning that he saw their feeding and polypide excursions.¹² Leendert Bomme encountered cilia on the tentacles in freshwater as well as marine bryozoans and connected 'a strong whirlpool, sucking all nearby objects to the animal into its stomach' with the ciliary activity 'permanent movement from down upwards, across the tip, and downwards again'.¹³ It should be noticed that Bomme was the first to record the travel of the ciliary movements over the tentacle tips in phylactolaemates. He criticized the inferences of Trembley,¹⁴ but since Bomme's brief report was apparently little known at that time, some later investigators neglected it and followed Trembley's opinion (for example, Lazzaro Spallanzani).¹⁵ Filippo Cavolini¹⁶ was criticized by Lucien Joliet¹⁷ for attributing the water currents to permanent tentacle movements, mainly because he did not recognize the tentacular cilia. However, precise examination of Cavolini's text shows that though he connected the lophophore with the origination of currents and often mentioned that 'tentacoli perpetuamente oscillano', he never wrote that their movements created these currents.¹⁸ Cavolini gave the most detailed descriptions of marine bryozoan feeding in the 18th century, recording some behavioural reactions (pharyngeal rejection, single expansion of lophophore, bending of a single or 2-3 tentacles deep inside the lophophore, waving tentacular movements, bringing straight tentacles together) for the first time.

3. Nineteenth century studies

After Rösel von Rosenhof's book mentioned above,¹⁹ several works on freshwater bryozoans with brief descriptions of their feeding behaviour were published during the next fifty years (summarized by William Sharpey²⁰ and George Allman²¹), and tentacular

ciliature was redescribed by Johann Eichhorn.²² Johann Steinbuch distinguished the cilia and their motion as the cause of water movements in *Plumatella repens* (Linnaeus) (as *Tubularia*), but, observing frequent retractions of polypides caused by an addition of extraneous particles, he inferred that the currents served chiefly, if not solely, for respiration.²³ Also, Steinbuch described and illustrated what seems to be water refiltration in the form of circular currents that had been already mentioned by Baker, who compared these currents with ‘the turning of a mill-wheel’.²⁴

As is evident from the old published descriptions and illustrations, living marine as well as freshwater bryozoans were often observed by investigators in the XIX century (Thompson,²⁵ de Blainville,²⁶ Gervais,²⁷ Nordmann,²⁸ Hassall,²⁹ Van Beneden,³⁰ Reid,³¹ Hancock,³² Leidy³³⁻³⁷ Gosse,³⁸ Busk,³⁹ Huxley,⁴⁰ Hincks,⁴¹ Müller,⁴² Smitt,⁴³,⁴⁴ Reichert,⁴⁵ Claparède,⁴⁶ Repiachoff,^{47,48} Ehlers,⁴⁹ Joliet,⁵⁰ Vigelius,⁵¹ Ostroumoff,⁵² Pergens,⁵³ Prouho).⁵⁴ Following Bomme,⁵⁵ many observers thought that effective strokes of lateral cilia were directed upwards along one side of a tentacle and downwards along the other (Grant,⁵⁶ Lister,⁵⁷ Dalyell,⁵⁸ Allman,⁵⁹ Hincks).⁶⁰ Based on this opinion, some authors concluded that there were ascending and descending water currents along consistent tentacle surfaces (for instance, John Fleming,^{61,62} and George Johnston).^{63,64} In contrast, Alpheus Hyatt was sure that cilia were ‘constantly vibrating towards the centre of the crown’, observing apparently their beating across the tentacle.⁶⁵ Similar statements were made by Heinrich Nitsche⁶⁶ and Karl Kraepelin⁶⁷ (see also below). Asajiro Oka apparently observed refiltering currents and thought that the cilia of the outer (abfrontal) tentacle surface generated the ascending current and those on the oral (frontal) surface produced the descending current.⁶⁸ The last four authors all worked with phylactolaemates.

Some investigators did not find the tentacular cilia, apparently due to the imperfection of microscopes. Noteworthy, Barthelemy Dumortier, who published one of the best anatomical descriptions of fresh-water bryozoans at that time, could not distinguish cilia on the tentacles in some cases (see also papers of François Raspail)^{69, 70} and made supposed absence of cilia the principal character of his new genus *Lophopus*.⁷¹

Robert Grant summarized previous works and gave a more detailed and correct description of marine bryozoan feeding. He was apparently the first to describe single and group tentacle flicks as a result of particle impacts in gymnolaemates.⁷² John Lister, who also worked with marine bryozoans, was the first to describe transport of particles along the frontal tentacle surface, their concentration in the pharynx before swallowing or rejection, and single and paired ‘hairs’ (setae) on the abfrontal tentacular surface. Lister’s illustrations show that he also found the groups of rigid cilia on the tips of the tentacles. Moreover, he was one of the first researchers who worked with living cyclostomates.⁷³

The most prominent investigation of this period on the feeding and behaviour of marine bryozoans was made by Arthur Farre.⁷⁴ He gave an excellent description of feeding in

seven ctenostomate and cheilostomate species. He was one of the first to use particles of carmine in his experiments to detect feeding currents. Many structures and reactions were recorded and beautifully illustrated by him for the first time: obliquely-truncate lophophores, scanning activity, testing-position, different variants of polypide protrusion, retraction and tentacle movements (for instance, writhing, curling, flattening of the lophophore when tentacles are almost perpendicular to the introvert), cessation of ciliature beating, etc. Moreover, he was the first to use the word 'vortex' which became very popular later to describe water turbulence within and nearby the lophophore (Bomme,⁷⁵ for instance, had used the Dutch word 'voortbrengen' = whirlpool) and proposed the filtration mechanism of bryozoan feeding. After Trembley,⁷⁶ Baker^{77, 78} and Bomme's⁷⁹ observations it was generally accepted that the feeding process included a directed water transference into the tentacle crown by ciliary activity as well as the activity of tentacles which 'assist the ciliary currents, by their varied movements, in obtaining food'.⁸⁰ Some observers believed that the tentacles 'catch' the food,⁸¹ whereas others were convinced that they 'arrest [retain] the prey' to 'conduct it to the mouth'.⁸² Despite the fact that Farre described various tentacle movements too, he thought they worked like 'a sieve, of a degree of fineness proportioned to the number of the arms and the consequent width of their intermediate spaces'.⁸³ According to his opinion, the size of filtered particles is regulated by the size of the mouth and the gaps between the tentacles. The former prohibits the swallowing of too large particles, and the latter, forming the sieve, lets small particles pass while retaining those of an intermediate size. Also Farre mentioned particle selection inside the pharynx resulting in swallowing or rejection.

John Dalyell gave very accurate and detailed descriptions of the behaviour in about 30 gymnolaemate and 4 phylactolaemate species.⁸⁴ He recorded many reactions, some of which were not described before. Following Arthur Hassall,⁸⁵ who pictured a bending of all the tentacles of the lophophore towards the mouth, Dalyell described and illustrated this reaction. In fresh-water bryozoans he recorded and described for the first time the pharyngeal rejection, concentration of particles in the pharynx before swallowing, and faster ciliature beat as a result of an increase of particle concentration. Dalyell compared feeding in bryozoans with that in sedentary polychaetes, thinking they both were capable of the 'absorption of muddy matter' and fed on 'small animals' only incidentally.⁸⁶ He was likely one of the first to perform experiments with bryozoans under light and low temperature influences.

George Allman observed feeding in about 20 species of phylactolaemates, and summarized much research on this topic in his monograph.⁸⁷ In *Fredericella sultana* (Blumenbach) he also illustrated the bending together only of tips of the tentacles for the first time.⁸⁸

Heinrich Nitsche encountered frontal cilia in sections of the tentacles of the freshwater bryozoan *Plumatella fungosa* (Pallas) (as *Alcyonella*).⁸⁹ He made his own observations

on living colonies and was able to distinguish ciliary beating along the frontal surface of the tentacles.⁹⁰ In his opinion, the entire lateral ciliature performed an ascending-descending movement (see above), but like Hyatt⁹¹ and Kraepelin,⁹² Nitsche thought that the cilia beat across the longitudinal axis of the tentacle. He first mentioned phase differences in the beating of the cilia, meaning a metachronal wave, and he found long rigid cilia both on frontal and abfrontal sides of the tentacles in the phylactolaemate bryozoan studied.⁹³ Samuel Silbermann summarized the existing data on presumed sensory structures in bryozoans. He described and illustrated paired and non-paired setae on the outer (abfrontal) side of tentacles in *Alcyonidium mytili* Dalyell.⁹⁴ He also encountered the fusiform cells with stiff cilia between ciliated cells on the inner side of the tentacle, but it is difficult to judge from his description and illustrations⁹⁵ whether or not these were the latero-frontal cells later described by Geneviève Lutaud⁹⁶ in cheilostomates (see below). In phylactolaemates similar fusiform 'sensory' cells with a hairy terminal process were described later by Adolf Gerwerzhagen⁹⁷ in *Cristatella mucedo* Cuvier and by Paul Brien⁹⁸ in *C. mucedo* and *Plumatella fungosa*. Gerwerzhagen reported that the cells found were irregularly distributed in bunches in the epithelial layer of the tentacle, whereas Brien believed that they are located on either side of a group of frontal cells with motile cilia, or even among them.^{99, 100}

Alpheus Hyatt described feeding in the fresh-water bryozoan *Fredericella sultana*.¹⁰¹ He was the first to report 'cagemaking',¹⁰² bending of tentacles to place a particle directly to the mouth, and rejection of the particles away by tentacles. Hyatt stressed that tentacles are not only feeding, but also sensory organs. Thomas Hincks summarized the existing data on bryozoan behaviour, and also made many of his own observations.¹⁰³ He described, in particular, some different lophophore shapes and tried to apply them to the current classification. Louis Calvet, using sections, found frontal cilia on the tentacles of some gymnolaemates. He believed that lateral cilia beat from outside towards the centre of the lophophore and back, across the tentacle and simultaneously from the top to its base. According to him, this beating created a turbulent water current towards the mouth and 'préhension' of particles, and the polypide then retracted, swallowing them.¹⁰⁴ Similar phenomenon was first described by Baker,¹⁰⁵ but Calvet thought that each swallowing was accompanied by retraction.¹⁰⁶ Calvet was also the first to mention repeated expansions and contractions of lophophore with straight tentacles, resembling pulsation.

4. Twentieth century studies

In the first half of the twentieth century, the most important investigations of marine bryozoan feeding and behaviour were made by Folke Borg. In 1923 he published a preliminary report of his research on bryozoan ciliary feeding,¹⁰⁷ and then gave a detailed analysis of the processes, with the description of many reactions.¹⁰⁸ Borg was the first to ascertain an oblique beating of the lateral cilia in respect to longitudinal tentacle axis, with effective strokes towards the outside. He compared the activity of these cilia, which differ

in phase, with the movement of an undulating membrane and thought that such undulation passes through the tentacle tips, as well as from one tentacle base to another. Borg correctly described the directions of water currents inside the lophophore and assigned an important role to the change of tentacle position and cilia beat direction as a reason for hydrodynamical changes inside and around the lophophore connected with feeding and particle rejection. In his opinion, the strongest current 'arises in the medial line of the tentacular crown in a direction straight downwards towards the mouth'¹⁰⁹ and 'mostly only those [particles which are contained in the current] in or near the medial line ... reach the mouth'.¹¹⁰ The particles are then ingested through action of the stomodaeum musculature that 'acts as a suction pump',^{111, 112} assisted by the pharyngeal ciliature.¹¹³ Borg recorded reversal, cessation, and deceleration of ciliary beating. Also he noted that the frontal cilia of tentacles played a subordinate role in the feeding process. Frontal cilia, beating from above downwards, rarely caused a slow transference of particles along the frontal surface towards the mouth. The mechanism of ciliary feeding described is enhanced by tentacular activities such as bending of individual tentacles directing particles towards the central current and lophophore contractions. Borg stressed that bryozoans perceived the nature of suspended particles and avoided ingesting inedible ones. In contrast, no food preferences were noted and particles were swallowed without apparent selection in experiments of Erich Ries.^{114, 115} For the first time Borg recorded a single lophophore contraction looking like a 'clap', brief series of tentacular pulsating movements during a rejection of particles, and bringing together only tips of the tentacles (accompanying a particle rejection in some cases).¹¹⁶

Ernst Marcus made many observations on marine as well as freshwater bryozoans,¹¹⁷⁻¹²¹ but his main aim was experimental work with changes of temperature, salinity and mechanical stimuli,¹²² and he did not study their feeding behaviour specifically. Marcus believed that the frontal cilia were immovable and stiff in at least one ctenostomate, *Farrella repens* (Farre).¹²³ Chandler Brooks was probably the first to describe scanning behaviour in Phylactolaemata.¹²⁴

Daphne Atkins compared the tentacle ciliary activity and the water currents created in Entoprocta and marine Ectoprocta.¹²⁵ She was the first to use the term 'metachronal wave' in describing the beating of the lateral cilia, and studied the operation of the ventral rejection tract in the ctenostomate *Flustrellidra hispida* (Fabricius) first described by Henry Proucho.¹²⁶ In *Pherusella tubulosa* (Ellis and Solander) (as *Pherusa*),^{127, 128} Atkins also divided the reactions connected with feeding behaviour into three groups that serve to: (1) increase the number of food particles, (2) prevent unwanted particles from reaching the mouth, (3) reject useless particles which have already reached the region of the mouth. Following Borg,¹²⁹ she suggested an important role of the frontal cilia situated at the base of tentacles in forming and directing water currents to the mouth.

In spite of two centuries of intensive observations, Carl Cori gave credit to the frontal

cilia in creating the main water current.¹³⁰ He thought that their beat was across the tentacle whereas lateral cilia were motionless or beat incidentally. This conclusion was strange because Cori saw the ascending-descending lateral ciliary movements, and described and illustrated them in some ctenostomates, comparing them with clock-wise ciliary movements in Kamptozoa and Phoronis.¹³¹ Using his own observations on the ctenostomate *Zoobotryon verticillatum* (Delle Chiaje) (as *Z. pellucidum*) and the data of Heinz Graupner, who investigated the anatomy of gymnolaemate tentacles,¹³² Cori suggested that the immobile lateral cilia formed a filter, closing narrow intertentacular gaps between the bases of all tentacles.¹³³ Sucking action of the frontal cilia resulted in water transference through the filter where suspended particles were retained, and, becoming associated with the mucus strands, conveyed to the pharynx. Like Lister,¹³⁴ Cori illustrated the stiff apical tentacle setae in *Z. verticillatum* without mentioning them in the text.¹³⁵ He also recorded the pushing away of particles by tips of tentacles for the first time.

Lars Silén investigated the functioning of the digestive tract of gymnolaemates in detail.¹³⁶ According to him the delivery of captured particles to the mouth proper is mainly performed by ciliary beating in the area around the mouth, stomodaeum and neighbouring part of the pharynx. Periodical contractions of the mouth may assist this transfer and take place independently from pharyngeal dilations and contractions resulting in food suction and swallowing.^{137,138} Later Silén described some zooidal behavioural reactions involving tentacles and entire polypides, and serving oviposition¹³⁹ and sperm release¹⁴⁰ (see also papers of Marcus,¹⁴¹ Bullivant,¹⁴² Cook,¹⁴³ Gordon,¹⁴⁴ Ström,¹⁴⁵ Chimonides and Cook,¹⁴⁶ Dyrinda and King,¹⁴⁷ summarized in the paper of Shunatova and Ostrovsky).¹⁴⁸

Despite numerous studies there was no clear concept of a bryozoan feeding mechanism. Dividing suspension feeders into groups Bernhard Werner marked out 'filterers' and 'whirlers' in his paper.¹⁴⁹ In the latter group, cilia generate a current and water with suspended particles is transferred along the collecting surface where the particles are retained. Bryozoans were considered to belong to a group of 'non-filtering suspension feeders'. In good agreement with Cori's suggestion,¹⁵⁰ Carl Jørgensen speculated that mucus played an important role in particle retention.¹⁵¹ In contrast, John Bullivant suggested that in all lophophorates a sharp deflection of water current passing through the lophophore causes particles to be thrown towards the mouth in a way which he compared with impingement separation used in industrial processes, and claimed that mucus plays no part in the feeding of marine bryozoans.^{152,153} He also supposed that laterofrontal cilia found by Lutaud¹⁵⁴ on the tentacles of the cheilostomate *Pentapora foliacea* (Ellis and Solander) (as *Hippodiplosia*)^{155,156} may either detect suspended particles in the water and initiate polypide retraction, tentacle flicking or reversal of lateral cilia, with the aim of rejection, or act as a sieve which retain the particles. Bullivant found and measured the laterofrontal cilia in the cyclostomate *Disporella separata* Osburn for the first time and used the term 'laeoplectic' to describe metachronal waves formed during lateral cilia beating.^{157,158,159} Following Bomme¹⁶⁰ and Marcus,¹⁶¹ he noted that metachronal waves

transmitted over the tips of tentacles in phylactolaemates, but he doubted that this took place in marine bryozoans. In *Plumatella repens*, Bullivant recorded the abfrontal cilia and described particle rejection along the frontal ciliary tract on the tentacles. He also confirmed¹⁶² observations of Günther Becker who described the rejection of particles by ciliature reversal of all of the tentacles in marine bryozoans.¹⁶³

Dennis Gordon investigated tentacular ciliature in the gymnolaemate *Cryptosula pallasiana* (Moll) and found that laterofrontal cilia performed occasional flicking movements.¹⁶⁴ He also described abfrontal cilia of two types that alternate with each other - short tufts of several immobile cilia and solitary cilia, and found the 'tuft of cilia at the tentacle tip'.¹⁶⁵

Bullivant's hypothesis was met with enthusiasm by some researchers, and John Ryland called bryozoans 'impingement feeders'.¹⁶⁶ Thomas Gilmour proposed a more detailed explanation of lophophorate feeding on the basis of the hypothesis in question and some theoretical generalizations.^{167, 168} He considered the process as 'simultaneous acceptance of food particles and rejection of inedible wastes probably' depending 'on the impingement or inertial impaction of heavy, potentially inedible particles on rejectory surfaces and the filtration or direct interception of light, potentially edible particles by cilia' on the lophophore.¹⁶⁹ In his opinion light particles are either thrown directly to the mouth with the incoming current or, if they move with a deflecting component of the current, are caught by the oral area cilia in gymnolaemates. Heavy particles are collected on the frontal surfaces by impingement and conveyed by the reversal of their cilia beating towards the tentacle tips. Following Bullivant,¹⁷⁰ Gilmour described the particle rejection along the frontal surface of the tentacles in the fresh-water bryozoan *Plumatella* sp., and also found this type of rejection in *Membranipora* sp. Gilmour also described the laterofrontal cilia on the tentacles of both gymnolaemate and phylactolaemate species (see above), but he supposed they work as sensory organs only.

The impingement model of lophophorate feeding was strongly criticized by Richard Strathmann,^{171, 172, 173} who argued that the velocities and the particle sizes and densities are too low for feeding by inertial impaction to operate.^{174, 175, 176} Instead, Strathmann speculated that suspended particles are captured by a combination of local reversal of the lateral cilia, which produces a local reversal of the water current, and inward flicking of the tentacles. The reversed current transfers the particles to the frontal side of the tentacle and helps to retain and transport them along the frontal surface towards the mouth (so-called up-stream collecting system according to terminology of Claus Nielsen and Jørgen Rostgaard).^{177, 178} It was suggested that this reversal must be induced by particles contacting either the lateral cilia themselves or the laterofrontal cilia. The latter may act as a sieve at the same time. This mechanism was first proposed for echinoderm larvae¹⁷⁹ and then extrapolated to lophophorates,^{180, 181} and Michael Hart presented a possible physiological explanation for its existence.¹⁸² In the ctenostomate *Flustrellidra hispida*,

Strathmann used high speed cinefilms to observe local alterations of metachrony of ciliary bands in places where the particles arrived. Though local reversal of the effective ciliary stroke could not be directly observed, these alterations were suggested to be the consequences of the reversals.¹⁸³ Strathmann did not report the transport of particles by frontal cilia in bryozoans, only that the transport along the frontal tentacle surface is aided by the reversal of the lateral cilia. He doubted that frontal cilia could transport the particles 'since only a few [of them] would be moving a particle against the action of many lateral cilia'.¹⁸⁴ Strathmann also was the first to measure the length of the lateral cilia in *F. hispida*,¹⁸⁵ to record frontal transport of particles by means of jumps and to investigate in detail flicking activity, which sometimes accompanied the proximal particle transport along the flicking tentacle.¹⁸⁶

Reactions to Strathmann's hypothesis were mixed. Some authors accepted it, but gave no direct evidence in support (Ryland,¹⁸⁷ Winston,^{188, 189} LaBarbera,¹⁹⁰ McKinney and Jackson,¹⁹¹ Hart,^{192, 193}). John Ryland tried to reach a compromise between the hypothesis of Bullivant (see above) and that of Strathmann.¹⁹⁴ He wrote that 'heavier particles may, as Bullivant suggested, be projected towards the mouth; but mostly it seems that particles are retained ... by local reversal of the lateral cilia'.¹⁹⁵ Ryland also discussed the significance of the width of the intertentacular gap and the length of lateral cilia in relation to Strathmann's opinions on particle capture.^{196, 196} According to Ryland's measurements of lophophores in different marine bryozoans, filtration should be fully effective in approximately the lower half of the tentacle crown since the lateral cilia should close the intertentacular gaps.^{198, 199} This idea was thereafter questioned and modified. James Markham and John Ryland²⁰⁰ and William Sanderson, John Thorpe and Andrew Clarke²⁰¹ found that the lateral cilia on the tentacles of *Flustrellidra hispida* and *Himantozoum antarcticum* (Calvet) correspondingly, 'exhibited a subterminal maximum length, decreasing proximally over more than half the tentacle and distally in about the last quarter'. Thus it was concluded that the actual closure of much of the intertentacular gap by the lateral cilia occurs along less than a quarter of the tentacle length or even lesser. The mechanism of local reversal would in principle be possible here as well, but it probably occurs in addition to other particle capture methods (see below). Filtration would be less effective in the upper part of the tentacle crown, where the distance between the tentacles is larger, but flicking might supplement the food collecting mechanism.

Michael Best and John Thorpe²⁰² denied the major role of local reversals in particle capture and transport towards the mouth.^{203, 204} Working on clearance rates and velocities of feeding currents in some gymnolaemates, they observed a frontal transport of particles downward and assumed that it might be a result of local reversals.²⁰⁵ At the same time Best and Thorpe came to the conclusion that the feeding process in the ctenostomate *Flustrellidra hispida* cannot be entirely explained by the local reversal-flicking hypothesis.²⁰⁶ They calculated, for instance, that the rate with which the pharynx is normally filled and the quantity of swallowed food particles would imply several flicks and more than 1000 local

ciliary reversals per second, which appears to be impossible. Following Borg²⁰⁷ and Atkins,²⁰⁸ Best and Thorpe supposed that long frontal cilia at the bases of the tentacles take an important part in bryozoan feeding. These cilia are used to direct the main feeding current towards the mouth, and 'the bulk of particles travel rapidly downwards in the current in the central area of the lophophore cone and proceed directly to the area above the mouth where they gather and are continuously ingested'.²⁰⁹ Local reversals and particle transport by frontal cilia are considered to be possible supplementary mechanisms.^{210,211} Best and Thorpe were the first to measure current velocities inside the lophophore.²¹²

The most exhaustive research on individual as well as group behaviour and associated structures was made by Judith Winston, who investigated about 80 species of marine bryozoans.²¹³⁻²¹⁶ She described and classified most of the feeding behavioural reactions known, recording many of them (collection of particles in a group inside the expanded lophophore near the mouth area, bending of all tentacle tips inside and outside the lophophore used for rejection at high particle concentrations, twisting activity, polypide retraction from under the particle, 'avoidance' retraction) for the first time. Winston also summarized the data on inferred sensory structures²¹⁷ and noted that the metachronal wave did not travel over the tentacle tips²¹⁸ (a fact which Bullivant doubted).²¹⁹ According to the prevailing reactions she distinguished 'filterers', 'tentacle feeders', 'scanners', 'cage-captors' and 'particle jugglers', and suggested that behavioural differences depend both on the physical and chemical characteristics of the food particles and behavioural characteristics of the various species.²²⁰ She also investigated parameters of the lophophore and its shape, showed similar trends in its symmetry within three main taxa of marine bryozoans and discussed their significance in relation to zooidal activity, colony structure (zooid arrangement and growth form) and environment. Winston argued that the complexity of zooidal behaviour was correlated with the level of colonial integration. The most complex individual behaviour was mainly found in poorly integrated colonies, and the most complex forms of group behaviour were discovered in highly integrated ones. The variety of reactions also often depended on polypide size: larger polypides perform a greater range of reactions involving different movements of tentacles and the entire polypide, whereas in small ones ciliary activity was the most important component of feeding behaviour as it was shown in cyclostomates and ctenostomates. Winston stressed that during particle rejection polypides could 'utilize a combination of methods depending on the size and concentration of the particles'.²²¹

Matthew Dick added to Winston's observations, recording for the first time in gymnolaemates an autocleaning of the colony surface by special activity of polypides in association with stiff abfrontal tentacular setae.²²² He described different types of rejection processes (particle flinging and partial retraction of polypide accompanying by pharyngeal rejection were described for the first time) and distinguished 'ejection' and 'rejection', and was the first to discuss the possible evolution of some behavioural

reactions in bryozoans. Following Winston, who suggested that abfrontal ciliary structures ‘could have some role in controlling polypide orientation relative to other polypides’,²²³ Dick speculated that ‘long abfrontal cilia’ perceive ‘tactile stimulation from the tentacles of surrounding polypides’ as well as ‘vectors’ of flow around the lophophore.²²⁴

Patricia Cook observed feeding behaviour of about 40 cheilostomates.²²⁵ In some species she described different variants of a lophophore protrusion and retraction in details, noting the specific sensitivity relative to the external irritants. Cook pictured several tentacle crown shapes,²²⁶ and described a temporary lophophore expansion with the tips of tentacles directed towards the colony surface for the first time. Since this reaction was “colony-wide”, the author suggested that it might be a colonial rejection, accompanied by ciliary reversal and connected with cleaning, sperm release or other unknown functions.

Ewan Hunter and Roger Hughes described an unusual variant of feeding in the cheilostomate *Celleporella hyalina* (Linnaeus) under high particle concentrations when ‘food particles were drawn from below the lophophore into the feeding crown by a current generated by the lateral cilia’.²²⁷ According to their observations ‘particles continued to move upwards and out of the lophophore, unless ciliary reversal occurred, bringing particles down’ to where finally they ‘were drawn into the mouth on a current generated by cilia within the pharynx’. A similar feeding method was earlier described by Borg.²²⁸ It is obvious from the texts that all of the authors postulate simultaneously changing ciliary movement on the tentacles, but not the local reversals proposed by Strathmann.^{229, 230} Hunter and Hughes also recorded a retaining of particles inside the lophophore when the tentacles cramed them into the mouth.²³¹

Hans Riisgård, Patricio Manríques²³² and Claus Nielsen²³³ reviewed a wide range of literature dealing mainly with physical aspects of particle capture by different suspension-feeders.²³⁴⁻²⁴⁵ Using theoretical considerations and non-contradictory ideas from conflicting theories of bryozoan feeding mechanism (see above) as well as their own data they proposed that the laterofrontal cilia form a mechanical filter which ‘strains the water while the central current, created by the special lophophore pump-design, and the action of flicking tentacles in co-operation clean the filter and transport the particles towards the mouth’.²⁴⁶ The laterofrontal cilia are considered to be multifunctional sensors that trigger tentacle flicking and local reversal of lateral cilia and serve as a mechanical sieve. By this means three main components of particle capture in bryozoans are effected: the direct transfer of the particles within the middle part of the central current to the mouth, particle impacts to a tentacle resulting in a downward transport by frontal cilia, and tentacle flicks conveying particles into the central lophophore current and triggered by the particle impacts with laterofrontal cilia. Flicking is stated to prevail in catching the particles in comparison with frontal transport,²⁴⁷ and the latter being impossible in at least some cyclostomates since frontal cilia are absent.^{248, 249} This is why every particle, that collides

with a tentacle, is captured by a single tentacle flick directing it into the central current in *Crisia eburnea* (Linnaeus). In all species investigated the frequency of flicking depends on concentration of particles. Riisgård and Manríques did not observe the capture of particles by local reversals and disturbances in ciliary metachrony and doubted compatibility of their proposed mechanism with Strathmann's model, but did not deny it entirely.²⁵⁰ They also noted that some results obtained by Best and Thorpe²⁵¹ and Sanderson and Thorpe²⁵² may be disputed because they had used too high concentrations of particles, tending to disturb normal feeding behaviour.²⁵³

Natalia Shunatova and Andrew Ostrovsky gave comparative descriptions of the behaviour in 40 species and subspecies of marine bryozoans.²⁵⁴ Some of the activities (cleaning of the colony surface using ascending water current created by reversal of the lateral cilia; cleaning of the colony surface by the inclined polypide; bending of the tentacle tips inside the lophophore during polypide protrusion; particle capture by tentacle expansion; pulsation of polypides bringing tentacles together within a lower part of the crown; particle rejection by bringing straight tentacles together; particle rejection in clumps by a series of repeated retractions and protrusions of polypide) were recorded for the first time. Behavioural reactions described were classified using a morphological approach based on the structure (tentacular ciliature, tentacles and entire polypide) performing the reaction. Authors discussed both normal and so-called spontaneous reactions as well as differences and similarities in individual autozooidal behaviour and their probable causes. They also suggested that in some species polypides can detect particles at some distance and that polypide protrusion and retraction might be the basis of the origin of some other individual activities. In marine bryozoans individual autozooidal behaviour was 'considered to be a flexible and sensitive system of reactions in which the activities can be performed in different combinations and successions, and can be switched depending on the situation'.²⁵⁵

Irina Antipenko studied in detail feeding of four fresh-water bryozoans recording many activities for the first time.²⁵⁶ For instance, testing-position was recorded for Phylactolaemata for the first time (in *Fredericella sultana*). She showed that, in contrast with marine bryozoans, frontal particle transport is one of the main feeding mechanisms (together with mechanical-filtering and tentacle flicking) in phylactolaemates. Small particles slide quickly whereas middle-sized ones usually move in jumps along the tentacle frontal surface toward the mouth. Instead, large particles are rejected being lifted by a reversal of frontal ciliature beat. When such a particle reaches a tentacle tip the rejecting tentacle sometimes sharply deflects outward shaking the particle off. If a particle is of 'intermediate' size, it often starts to travel down and up before it is either swallowed or rejected. Large particles can also be rejected by a brief reversal of lateral ciliature of all tentacles, expansion of lophophore, bending either of a single or several tentacles outside, sharp shaking of all of the tentacles. In *Cristatella mucedo* rejecting particles can be rolled from tentacle to tentacle towards the end of the lophophore 'arm'. The polypide can cross

'arms' in this species, and perform a 'clap' to catch more particles. Finally, Antipenko stressed the great similarity in autozooidal behaviour of *F. sultana* (possessing ring lophophore) and that of marine bryozoans.

Quantitative characteristics of feeding in marine bryozoans have been investigated since the works of John Bullivant²⁵⁷ and N. Ravindranatha Menon²⁵⁸ who measured clearance rates by calculating theoretical volumes of water cleared of particles per unit time, and Judith Dudley [Winston]²⁵⁹ and John Ryland²⁶⁰ who studied dimensional parameters of the lophophore.²⁶¹⁻²⁶⁶ Best and Thorpe,^{267, 268, 269} Sanderson *et al.*,²⁷⁰ Sanderson and Thorpe²⁷¹ and Riisgård and Manríques²⁷² made direct measurements of feeding current velocities for various marine Bryozoa. These studies indicate that velocities vary across the diameter of the lophophore and are the strongest axially.^{273, 274} Velocities of feeding current may also vary considerably depending on a number of parameters including food type, food particle size and concentration and the extent to which the colony has been starved.^{275, 276, 277} Both intraspecific and interspecific variations in lophophore behaviour and morphology were documented, although generally parameters of lophophore width, tentacle length and tentacle number correlate quite strongly.²⁷⁸⁻²⁸³ It was supposed that interspecific (and intergeneric) differences in behaviour and morphology of the lophophore probably result in corresponding differences in diet,²⁸⁴⁻²⁹² and 'that bryozoans [with] ... a substantial difference in the size of the lophophore and tentacle number ... may be able to coexist by means of niche partitioning'.^{293, 294} Phytoplankton (mainly, diatoms) is almost certainly the most important food source for shallow-water bryozoans, but various motile protists (like ciliates and flagellates), small planktonic arthropods, detritus, bacterial suspensions and dissolved organic matter are considered to be potential food sources as well.²⁹⁵⁻³⁰⁶ Seasonal and tidal changes in food composition and concentrations have been recorded³⁰⁷⁻³¹⁰ and intraspecific behavioural variability may be partially explained by these factors.³¹¹ Feeding is also influenced by an increased temperature.³¹²⁻³¹⁶

Current regimes have been shown to influence feeding of marine bryozoans. Relationships between ambient flow velocity, particle sizes and feeding success in the colonies of different growth forms were intensively studied by Beth Okamura.³¹⁷⁻³²⁰ She found that the capture of particles of different sizes may be affected in different ways under various conditions of flow and suggested a possible switch in feeding behaviour from filtering to capturing by tentacles.³²¹ It has been also shown that bryozoan feeding was significantly affected by the interacting influences of the flow velocity and degree of association with neighbouring suspension-feeders, for instance, other bryozoans,^{322, 323} and these conclusions conform with the results of previous work by Leo Buss³²⁴ and Michael Best and John Thorpe^{325, 326} on feeding-current interactions among neighbouring colonies.^{327, 328, 329} Feeding of some marine bryozoans was also investigated in laminar³³⁰ and turbulent flow.³³¹ Both studies showed that feeding rates increase with increasing flow velocity. Unfortunately, phylactolaemates are almost non-studied in these

respects.^{332, 333}

5. Conclusions

In conclusion, we would like to stress that in spite of more than 250 years of research our knowledge of bryozoan feeding behaviour is still far from satisfactory, and implied feeding mechanisms are based more on hypotheses and suggestions than on facts. Structural uniformity of the bryozoan polypide and tentacles implies common feeding mechanisms. It is generally accepted that bryozoans probably use several mechanisms depending on polypide morphology and size within different taxa, and changes of environment.³³⁴⁻³⁴⁰ Supposedly, the basic lophophore filter-feeding principle is added by tentacle flicking, frontal transport and, may be, local reversals. Some of these mechanisms can apparently be used simultaneously, but their importance for feeding is, probably, different. Further investigations are necessary to prove these speculations. Microscopical size and the impossibility to observe bryozoans in natural habitats are the main reasons of many existing problems and theoretical controversies. We hope that the complex approach of using sophisticated videoequipment, carefully planned and accomplished quantitative experiments and anatomical work^{341, 342, 343} could improve this situation.

6. Acknowledgements

We are greatly indebted to Professor Dr Claus Nielsen and Dr Mary E. Petersen, Zoological Museum, University of Copenhagen, and Mary Spencer Jones, The Natural History Museum, London, for kind help with literature. Sincere thanks are given to Dr Margit Jensen, Zoological Museum, University of Copenhagen, Dr Harry A. ten Hove, Institute for Systematics and Population Biology, Zoological Museum, University of Amsterdam, Ekaterina V. Lobastova, Biological Scientific Research Institute, St. Petersburg State University, Natalia V. Lentsman, Dept. of Invertebrate Zoology, St. Petersburg State University, Dr Elena A. Istomina, Seacamp Association, Florida, and Tatjana V. Ostrovskaja, Leningrad Zoo, for their assistance with translating. Our thanks to Drs William G. Sanderson, Michael A. Best and John P. Thorpe, Department of Environmental and Evolutionary Biology, University of Liverpool, Port Erin Marine Laboratory, for an unpublished review on bryozoan feeding. We are grateful to Dr Judith E. Winston, Virginia Museum of Natural History, Professor Dr Roger N. Hughes, School of Biological Sciences, University of Wales, Bangor, Dr Ken McKinney, Appalachian State University and Dr Paul D. Taylor, The Natural History Museum, for reading and criticizing earlier drafts of the manuscript, useful comments and much encouragement. Andrew N. Ostrovsky thanks the Danish Ministry of Education for a post-doctoral stipend which made the work in University of Copenhagen Library possible. The Otto Kinne Foundation is acknowledged for financial support.

Notes

- 1 N.N. Shunatova and A.N. Ostrovsky, 'Individual autozooidal behaviour and feeding in marine bryozoans', *Sarsia*, 86 (2001), 113-142.
- 2 N.N. Shunatova and A.N. Ostrovsky, 'Group autozooidal behaviour and chimneys in marine bryozoans', *Marine Biology*, 140 (2002), 503-518.
- 3 J.E. Winston, 'Polypide morphology and feeding behaviour in marine ectoprocts', *Bulletin of Marine Science*, 28(1) (1978), 1-31 (1).
- 4 A. Trembley, *Memoires pour servir a l'histoire d'un genre de polypes d'eau douce, a bras en forme de cornes*, 2 vols (Paris, Jean and Herman Verbeek, 1744), xvi, 324 (vol 2, 128-147, Plate 10, figures 8-9).
- 5 See for synonymy L.H. Hyman, *The invertebrates: smaller coelomate groups*, 5 (McGraw-Hill Book Company, New York, Toronto, London, 1959), 275-515.
- 6 H. Baker, *An attempt towards a natural history of the polype* (London, 1743), 248.
- 7 H. Baker, *Employment for the microscope* (London, 1753), xiv, 442.
- 8 Trembley, note 4.
- 9 Baker, note 7, p. 309.
- 10 See for further discussion Shunatova and Ostrovsky, note 1.
- 11 A.J. Rösel von Rosenhof, *Der monatlich-herausgegebenen Insecten-Belustigung*, 3 (Nürnberg, 1755), 624.
- 12 J. Ellis, *An essay towards a natural history of the corallines, and other marine productions of the like kind, commonly found on the coasts of Great Britain and Ireland* (London, 1755), 120.
- 13 L. Bomme, 'Bericht wegens een zonderling Zeeinsect, gevonden aan eenige Zeewieren, gevischt op het Strand van het Eiland', *Verhandelingen Uitgegeven door het Zeeuwisch Genootschap der Wetenschappen te Vlissingen*, 1 (1769), 394-402 (400).
- 14 Trembley, note 4.
- 15 L. Spallanzani, *Opusculi de physique, animale et végétale*, 2 vols (Geneve, 1777), vol 1, cxxiv, 255, vol 2, ii, 404.
- 16 F. Cavolini, *Memoire per servire alla storia de'polipi marini di Filippo Cavolini* (Napoli, 1785), 279.
- 17 L. Joliet, 'Contributions à l'histoire naturelle des Bryozoaires des côtes de France', *Archives de Zoologie expérimentale et générale*, 6 (1877), 193-304.
- 18 Cavolini, note 16, p. 244.
- 19 Rösel von Rosenhof, note 11.
- 20 W. Sharpey, 'Cilia', in *The cyclopaedia of anatomy and physiology*, edited by R.B. Todd (Sherwood, Gilbert and Piper, London, 1836), 606-638.
- 21 G.J. Allman, *A monograph of the fresh-water Polyzoa, including all the known species, both british and foreign* (Ray Society, London, 1856), 120.
- 22 J.C. Eichhorn, *Beiträge zur Naturgeschichte der kleinsten Wasserthiere in den Gewässern und um Danzig* (Danzig, 1776), 94.
- 23 J.G. Steinbuch, *Analecten neuer Beobachtungen und Untersuchungen für die Naturkunde* (Fürth, 1802), 135.
- 24 Baker, note 6, p. 22.
- 25 J.V. Thompson, 'On Polyzoa, a new animal discovered as an inhabitant of some zoophytes with a description of the newly instituted genera of *Pedicellaria* and *Vesicularia* and their species', *Zoological researches and illustrations*, 4(5) (1830), 89-102.

- 26 H.M.D. de Blainville, *Manuel d'Actinologie ou de Zoophytologie*, 2 vols (Paris, F. G. Levrault, 1834), 694.
- 27 M.P. Gervais, 'Recherches sur les Polypes d'eau douce des genres *Plumatella*, *Cristatella* et *Paludicella*', *Annales des Sciences Naturelles*, 2 Serie, 7 (Masson, Paris, 1837), 74-93.
- 28 M.A. Nordmann, 'Recherches microscopiques sur l'anatomie et le développement du *Tendra zostericola*, espèce de polype de la section des Bryozoaires', *Annales des Sciences Naturelles*, 2 Serie, 11 (Masson, Paris, 1839), 185-191.
- 29 A.H. Hassall, 'Supplement to a catalogue of Irish zoophytes', *The Annals and Magazine of Natural History*, 7(44, 45) (1841), 276-87, 363-373.
- 30 P.J. van Beneden, 'Recherches sur l'organisation des *Laguncula* et l'histoire naturelle des différents polypes Bryozoaires qui habitent la côte d'Ostende', *Nouveaux Mémoires de l'Académie de Bruxelles*, 18 (1844), 1-29.
- 31 J. Reid, 'Anatomical and physiological observations on some Zoophytes', *The Annals and Magazine of Natural History*, 16 (1845), 385-400.
- 32 A. Hancock, 'On the anatomy of the freshwater Bryozoa, with descriptions of three new species', *Annals and Magazine of Natural History*, 5 (1850), 173-204.
- 33 J. Leidy, 'On the genus *Plumatella*', *Proceedings of the Academy of Natural Sciences of Philadelphia*, 5 (1850-1851), 261-262.
- 34 J. Leidy, 'On *Cristatella*', *Proceedings of the Academy of Natural Sciences of Philadelphia*, 5 (1850-1851), 265-266.
- 35 J. Leidy, 'On some American fresh-water Polyzoa', *Proceedings of the Academy of Natural Sciences of Philadelphia*, 5 (1850-1851), 320-322.
- 36 J. Leidy, 'On *Urnatella gracilis* and a new species of *Plumatella*', *Proceedings of the Academy of Natural Sciences of Philadelphia*, 7 (1854-1855), 191-192.
- 37 J. Leidy, 'Remarks on Polyzoa', *Proceedings of the Academy of Natural Sciences of Philadelphia*, 10 (1858), 188-190.
- 38 P.H. Gosse, 'Notes on some new or little-known marine animals', *The Annals and Magazine of Natural History*, 2 Serie, 16 (1855), 27-36.
- 39 G. Busk, 'Zoophytology', *Quarterly Journal of Microscopical Science*, 4 (1856), 308-312.
- 40 T.H. Huxley, 'Note on the reproductive organs of the cheilostome Polyzoa', *Quarterly Journal of Microscopical Science*, 4 (1856), 191-192.
- 41 T. Hincks, 'Note on the ovicells of the cheilostomatous Polyzoa', *Quarterly Journal of Microscopical Science*, New Series, 1 (1861), 278-281.
- 42 F. Müller, 'On the common nervous system (Kolonialnervensystem) of the Bryozoa (Polyzoa), exemplified in *Serialaria Coutinhii*, n. sp.', *Quarterly Journal of Microscopical Science*, New Series, 1 (1861), 300-305.
- 43 F.A. Smitt, 'Om Hafs-Bryozoernas utveckling och fettkroppar', *Öfversigt af Kongliga Vetenskapsakademien Förhandling*, 1 (1865), 5-50.
- 44 F.A. Smitt, 'Kritisk fürteckning öfver Skandinaviens Hafs-Bryozoer', *Öfversigt af Kongliga Vetenskapsakademien Förhandling*, 2 (1865), 115-533.
- 45 H. Reichert, 'Vergleichende anatomische Untersuchungen über *Zoobotryon pellucidus* (Ehrenberg)', *Abhandlungen der königlichen Akademie der Wissenschaften zu Berlin*, 2 (1869), 233-338.
- 46 E. Claparède, 'Beiträge zur Anatomie und Entwicklungsgeschichte der Seebryozoen', *Zeitschrift für Wissenschaftliche Zoologie*, 21 (1871), 138-174.

- 47 W. Repiachoff, 'Zur Entwicklungsgeschichte der *Tendra zostericola*', *Zeitschrift für wissenschaftliche Zoologie*, 25 (1875), 129-142.
- 48 W. Repiachoff, 'Zur Naturgeschichte der chilostomen Bryozoen', *Zeitschrift für wissenschaftliche Zoologie*, 26 (1876), 139-160.
- 49 E. Ehlers, '*Hypophorella expansa*. Ein Beitrag zur Kenntniss der minirenden Bryozoen', *Abhandlungen der physikalischen Classe der königlichen Gesellschaft der Wissenschaften zu Göttingen*, 21 (1876), 3-157.
- 50 Joliet, note 17.
- 51 W.J. Vigelius, 'Morphologische Untersuchungen über *Flustra membranaceo-truncata* Smitt', *Biologisches Centrblatt*, 23 (1884), 705-721.
- 52 A. Ostroumoff, 'Die Bryozoen der Bucht von Sebastopol. Vollständigere Ausgabe mit einem ganz neuen Theile über die Morphologie der Bryozoen', *Proceedings of the Society of Naturalists of Kazan*, 16(2) (1886), 1-124 (in Russian).
- 53 E. Pergens, 'Untersuchungen an Seebryozoen', *Zoologischer Anzeiger*, 12 (1889), 504-510, 526-533.
- 54 H. Proucho, 'Contribution à l'histoire des Bryozoaires', *Archives de Zoologie expérimentale et générale*, 10 (1892), 557-656.
- 55 Bomme, note 13.
- 56 R.E. Grant, 'Observations on the structure and nature of *Flustra*', *Edinburgh New Philosophical Journal*, 3 (1827), 107-118, 337-342.
- 57 J.J. Lister, 'Some observations on the structure and functions of tubular and cellular polypi, and of Ascidiæ', *Philosophical Transactions of the Royal Society of London*, 1 (1834), 365-388.
- 58 J.G. Dalyell, *Rare and remarkable animals of Scotland, represented from living subjects: with practical observations on their nature*, 2 vols (Van Voorst, London, 1847-1848), vol 1, xii, 268, vol 2, iv, 322.
- 59 Allman, note 21.
- 60 T. Hincks, '*A history of the British marine Polyzoa*', 2 vols (Van Voorst, London, 1880), 601.
- 61 J. Fleming, 'Observations on the *Sertularia cuscuta* Ellis', *Memoirs of the Wernerian Natural History Society*, 4(2) (1823), 485-491.
- 62 J. Fleming, '*A history of British animals*', (Edinburgh, London, 1828), xxiii, 565.
- 63 G. Johnston, '*A history of the British zoophytes*', (Edinburgh, London, Dublin, 1838), 341.
- 64 G. Johnston, '*A history of the British zoophytes*', 2nd edition, (Van Voorst, London, 1847), xiv, 488.
- 65 A. Hyatt, 'Observations on Polyzoa. Suborder Phylactolaemata', *Proceedings of the Essex Institute*, vols 4 and 5, (Salem, 1866-1868), iv, 103 (9).
- 66 H. Nitsche, *Beiträge zur Anatomie und Entwicklungsgeschichte der phylactolaemen Süßwasserbryozoen, insbesondere von Alcyonella fungosa* Pall. Akademisch Dissertation, (Grimm and Maass, Berlin, 1868), 57.
- 67 K. Kraepelin, 'Die Deutschen Süßwasser-Bryozoen. Anatomisch-systematischer Teil', *Abhandlungen aus dem Gebiete der Naturwissenschaft herausgeben von dem Naturwissenschaftlicher Verein in Hamburg*, 10 (1887), 1-168.
- 68 A. Oka, 'Observations on fresh-water Polyzoa. (*Pectinatella gelatinosa*, nov. sp.)', *Journal of the College of Science, Imperial University of Japan, Tokyo*, 4 (1891), 89-150.
- 69 F.V. Raspail, 'Analyse du polypier de l'*Alcyonelle* des etangs Ferussac', *Bulletin des Sciences Mathématique*, 9 (1828), 138-141.

- 70 F.V. Raspail, 'Histoire naturelle de l'*Alcyonella fluviatile* (*A. stagnorum*, Lamk.) et de tous les genres voisins, consideres, soit sous le rapport de leur organisation et de leur identite specifique, sout sous le rapport physiologique de leurs tentacules avec les branchies des mollusques et des animalcules ou infusoires ou sermatiques', *Memoires de la Societe d'Histoire Naturelle de Paris*, 4 (1828), 75-165, 246.
- 71 B.-C. Dumortier, 'Recherches sur l'anatomie et la physiologie des polypiers composés d'eau douce', *Bulletins de l'Académie Royale des Sciences, Bruxelles*, 2 (1835), 421-454.
- 72 Grant, note 56.
- 73 Lister, note 57, p. 384, plate XII, figures 2, 5.
- 74 A.B. Farre, 'Observations on the minute structure of some of the higher forms of polypi, with views of a more natural arrangement of the class', *Philosophical Transactions of the Royal Society of London*, 1 (1837), 387-426.
- 75 Bomme, note 13.
- 76 Trembley, note 4.
- 77 Baker, note 6.
- 78 Baker, note 7.
- 79 Bomme, note 13.
- 80 Hincks, note 60, volume 1, p. xv.
- 81 Baker, note 6, p. 23.
- 82 Johnston, note 64, p. 258.
- 83 Farre, note 74, p. 408.
- 84 Dalyell, note 58.
- 85 Hassall, note 29, plate VI, figure 1.
- 86 Dalyell, note 58, volume 2, p. 281.
- 87 Allman, note 21.
- 88 Allman, note 21, Plate IX, figure 2.
- 89 Nitsche, note 66.
- 90 See also Kraepelin, note 67.
- 91 Hyatt, note 65, p. 9.
- 92 Kraepelin, note 67.
- 93 Nitsche, note 66, p. 27, plate III, figure 31.
- 94 S. Silbermann, 'Untersuchungen über den feineren Bau von *Alcyonidium mytili*', *Archiv für Naturgeschichte*, 72 (1906), 265-308.
- 95 Silbermann, note 94, 277-8, plate XIX, figure 13A-C.
- 96 G. Lutaud, 'Sur la ciliature du tentacule chez les Bryozoaires Chilostomes', *Archives de Zoologie expérimentale et générale (Notes et Revue)*, 92 (1955), 13-19.
- 97 A. Gerwerzhagen, 'Beiträge zur Kenntnis der Bryozoen. I. Das Nervensystem von *Cristatella mucedo* Cuv.', *Zeitschrift für Wissenschaftliche Zoologie*, 107 (1913), 309-345.
- 98 P. Brien, 'Classe des Bryozoaires', in *Traité de Zoologie*, edited by P.-P. Grasse, 5(2) (Masson et Sie, Paris, 1960), 1053-1335.
- 99 For further discussion see G. Lutaud, 'The bryozoan nervous system', in *The Biology of Bryozoans*, edited by R.M. Woollacott and R.L. Zimmer (Academic Press, New York, London, 1977), 377-410.
- 100 H. Mukai, K. Terakado and C.G. Reed, 'Bryozoa', in *Microscopic Anatomy of Invertebrates*, edited by F.W. Harrison, vol 13, (New York, Wiley-Liss, 1997), 45-206.
- 101 Hyatt, note 65, p. 9.

- 102 For terminology see J.E. Winston, note 3.
- 103 Hincks, note 60.
- 104 L. Calvet, 'Contribution à l'histoire naturelle des Bryozoaires Ectoproctes marins', *Travailes de l'Institut de Zoologie de l'Université de Montpellier*, Nouvelle Série, 8 (1900), 1-488 (4).
- 105 Baker, note 7.
- 106 Criticized in F. Borg, 'Studies on recent cyclostomatous Bryozoa', *Zoologiska Bidrag från Uppsala*, 10 (1926), 181-250.
- 107 F. Borg, 'On the structure of cyclostomatous Bryozoa', *Arkiv für Zoologie*, 15 (1923), 1-17 (8).
- 108 Borg, note 106.
- 109 Borg, note 106, p. 247.
- 110 Borg, note 106, p. 248.
- 111 See also F. Braem, 'Victorella sibogae Harmer', *Zeitschrift für Morphologie und Ökologie der Tiere*, 36 (1940), 267-78.
- 112 L. Silén, 'On the division and movements of the alimentary canal of the Bryozoa', *Arkiv für Zoologie utgivet av Kongliga Svenska Vetenskapsakademien* 35, 3(12) (1944), 1-40.
- 113 Borg, note 106, p. 247.
- 114 E. Ries, 'Fütterungsversuche bei *Zoobotryon* (Bryozoa)', *Zeitschrift für Vergleichende Physiologie*, 23 (1936), 64-99.
- 115 See also Shunatova and Ostrovsky, note 1.
- 116 Borg, note 106.
- 117 E. Marcus, 'Beobachtungen und Versuche an lebenden Meersbryozoen', *Zoologische Jahrbücher Abteilung für Systematik, Ökologie und Geographie der Tiere*, 52 (1926), 1-102.
- 118 E. Marcus, 'Beobachtungen und Versuche an lebenden Süßwasserbryozoen', *Zoologische Jahrbücher Abteilung für Systematik, Ökologie und Geographie der Tiere*, 52 (1926), 279-350.
- 119 E. Marcus, 'Bryozoa', *Die Tierwelt der Nord und Ostsee*, 4 (7c) (1926), 1-100.
- 120 E. Marcus, 'Über *Lophopus crystallinus* (Pall.)', *Zoologische Jahrbücher Abteilung für Anatomie und Ontogenie der Tiere*, 58 (1934), 501-606.
- 121 E. Marcus, 'Sobre Bryozoa do Brasil', *Boletins da Faculdade de Filosofia, Ciências e Letras. Universidade de São Paulo XXII*, *Zoologia*, 5 (1941), 3-208.
- 122 Summarized in S.F. Harmer, 'Recent work on Polyzoa', *Proceedings of the Linnean Society of London*, 143 (1931), 113-68.
- 123 E. Marcus, note 117.
- 124 C.M. Brooks, 'Notes on the statoblasts and polypids of *Pectinatella magnifica*', *Proceedings of the Academy of Natural Sciences of Philadelphia*, 81 (1929), 427-441.
- 125 D. Atkins, 'The ciliary feeding mechanism of the Entoproct Polyzoa, and a comparison with that of the Ectoproct Polyzoa', *Quarterly Journal of Microscopical Science*, New Series, 75(299) (1932), 393-423.
- 126 Proucho, note 54.
- 127 See also F. Braem, 'Über die Querstreifung im Pharynx der gymnolämen Bryozoen und über den Bau des Munddarms', *Zeitschrift für Morphologie und Ökologie der Tiere*, 36 (1940), 668-676.
- 128 Marcus, note 121.
- 129 Borg, note 106.

- 130 C.J. Cori, 'Ordnung der Tentaculata. Bryozoa', in *Handbuch der Zoologie* III, 2(5), (Walter de Gruyter and Company, Berlin, 1941), 263-374, 375-503.
- 131 Cori, note 130, p. 409, figures 349, 366, 492.
- 132 H. Graupner, 'Zur Kenntnis der feineren Anatomie der Bryozoen (Nervensystem, Muskulatur, Stützmembran)', *Zeitschrift für Wissenschaftliche Zoologie*, 136 (1930), 38-77.
- 133 For comparison see J.B. Markham and J.S. Ryland, 'Morphometry and function in the lophophore of *Flustrellidra hispida* (Bryozoa)', in *Bryozoaires Actuels et Fossiles: Bryozoa Living and Fossil*, edited by F.P. Bigey, *Bulletin de la Société des Sciences Naturelles de l'Ouest de la France*, Mémoire Hors Série, 1 (1991), 237-242.
- 134 Lister, note 57.
- 135 Cori, note 130, figure 346.
- 136 Silén, note 112.
- 137 See also M.A. Best and J.P. Thorpe, 'Effects of particle concentration on clearance rate and feeding current velocity in the marine bryozoan *Flustrellidra hispida*', *Marine Biology*, 77 (1983), 85-92.
- 138 E. Hunter and R.N. Hughes, 'The effect of cell concentration on colony growth and feeding in the bryozoan *Celleporella hyalina*', *Journal of the Marine Biological Association of the United Kingdom*, New Series, 73 (1993), 321-331.
- 139 L. Silén, 'The main features of the development of the ovum, embryo and oocidium in the oocioferous Bryozoa Gymnolaemata', *Arkiv für Zoologie*, 35A (1945), 1-34.
- 140 L. Silén, 'On the fertilization problem in the gymnolaematous Bryozoa', *Ophelia*, 3 (1966), 113-140.
- 141 Marcus, note 117.
- 142 J.S. Bullivant, 'Release of sperm by Bryozoa', *Ophelia*, 4 (1967), 139-142.
- 143 P.L. Cook, 'Observations on living Bryozoa', *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, 108 (1968), 155-160.
- 144 D.P. Gordon, 'Zooidal dimorphism in the polyzoan *Hippopodinella adpressa* (Busk)', *Nature (London)*, 219 (1968), 633-634.
- 145 R. Ström, 'Sexual reproduction in a stoloniferous bryozoan, *Triticella koreni* (G. O. Sars)', *Zoologiska Bidrag från Uppsala*, 38 (1969), 113-128.
- 146 P.J. Chimonides and P.L. Cook, 'Observations on living colonies of *Selenaria* (Bryozoa, Cheilostomata). II', *Cahiers de Biologie Marine*, 22 (1981), 207-219.
- 147 P.E.J. Dyrinda and P.E. King, 'Gametogenesis in placental and non-placental ovicellate cheilostome Bryozoa', *Journal of Zoology (London)*, 200 (1983), 471-942.
- 148 Shunatova and Ostrovsky, note 1.
- 149 B. Werner, 'Das Prinzip des andlosen Schleimfilters beim Nahrungswerb wirbelleser Meerestiere', *Internationale Revue der gesamten Hydrobiologie und Hydrographie*, 44 (1959), 181-215.
- 150 Cori, note 130.
- 151 C.B. Jørgensen, *Biology of suspension feeding*, (Pergamon Press, Oxford, London, New York, Paris, 1966), xv, 357.
- 152 J.S. Bullivant, 'The method of feeding of Lophophorates (Bryozoa, Phoronida, Brachiopoda)', *New Zealand Journal of Marine and Freshwater Research*, 2(1) (1968) 135-46.
- 153 J.S. Bullivant, 'A revised classification of suspension feeders', *Tuatara*, 16 (1969), 151-60.
- 154 Lutaud, note 96.
- 155 See also G. Lutaud, 'L'innervation du lophophore chez le Bryozoaire chilostome *Electra*

- pilosa* (L.)', *Zeitschrift für Zellforschung und mikroskopische Anatomie*, 140 (1973.), 217-234.
- 156 L.W. Smith, 'Ultrastructure of the tentacles of *Flustrellidra hispida* (Fabricius)', in *Living and Fossil Bryozoa*, edited by G.P. Larwood (Academic Press, London, New York, 1973), 335-42.
- 157 Bullivant, note 152, p. 137.
- 158 For general discussion see also M.A. Sleight and E. Aiello, 'The movement of water by cilia', *Acta Protozoologica*, 11 (1972), 265-277.
- 159 J.R. Blake and M.A. Sleight, 'Hydromechanical aspects of ciliary propulsion', in *Swimming and Flying in Nature*, edited by T.Y.T. Wu, C.J. Brokaw and C. Brennen (Plenum Press, New York, 1975), 1, 185-209.
- 160 Bomme, note 13.
- 161 Marcus, note 117.
- 162 J.S. Bullivant, 'The rate of feeding of the bryozoan, *Zoobotryon verticillatum*', *New Zealand Journal of Marine and Freshwater Research*, 2(1) (1968b), 111-133.
- 163 G. Becker, 'Untersuchungen über den Darm und die Verdauung von Kamptozoen, Bryozoen und Phoroniden', *Zeitschrift für Morphologie und Ökologie der Tiere*, 33 (1938), 72-127.
- 164 D.P. Gordon, 'Microarchitecture and function of lophophore in the bryozoan *Cryptosula pallasiana*', *Marine Biology*, 27 (1974), 147-163.
- 165 Gordon, note 164, p. 151.
- 166 J.S. Ryland, *Bryozoans* (Hutchinson University Library, London, 1970) 175 (43).
- 167 T.H.J. Gilmour, 'Ciliation and function of the food-collecting and waste-rejecting organs of lophophorates', *Canadian Journal of Zoology*, 56 (1978), 2142-2155.
- 168 D.I. Rubenstein and M.A.R. Koehl, 'The mechanisms of filter feeding: some theoretical considerations', *American Naturalist*, 111 (1977), 981-994.
- 169 Gilmour, note 167, p. 2152.
- 170 Bullivant, note 152.
- 171 R.R. Strathmann, 'Function of lateral cilia in suspension feeding of lophophorates (Brachiopoda, Phoronida, Ectoprocta)', *Marine Biology*, 23 (1973), 129-136.
- 172 R.R. Strathmann, 'Cinefilms of particle capture by an induced local change of beat of lateral cilia of a bryozoan', *Journal of Experimental Marine Biology and Ecology*, 62 (1982), 225-236.
- 173 R.R. Strathmann, 'Comment on Dr. Gilmour's views on feeding by hemichordates and lophophorates', *Canadian Journal of Zoology*, 60 (1982), 3466-3468.
- 174 For discussion see Rubenstein and Koehl, note 168.
- 175 S. Vogel, *Life in moving fluids: the physiological biology of flow* (Princeton University Press, Princeton, NJ, 1994), xiii, 467.
- 176 M.W. Hart and R.R. Strathmann, 'Mechanisms and rates of suspension feeding', in *Ecology of Marine Invertebrate Larvae*, edited by L. McEdward, Marine Science Series, (CRC Press, Boca Raton, FL, 1995), 193-221.
- 177 C. Nielsen and J. Rostgaard, 'Structure and function of an entoproct tentacle with a discussion of ciliary feeding types', *Ophelia*, 15 (1976), 115-140.
- 178 For review see also C. Nielsen, 'Structure and function of metazoan ciliary bands and their phylogenetic significance', *Acta Zoologica*, 68 (1987), 205-262.
- 179 R.R. Strathmann, 'The feeding behaviour of planctotrophic echinoderm larvae: mechanisms,

- regulation, and rates of suspension feeding', *Journal of Experimental Marine Biology and Ecology*, 6 (1971), 109-160.
- 180 R.R. Strathmann, T.L. Jahn and J.R.C. Fonseca, 'Suspension feeding by marine invertebrate larvae: clearance of particles by ciliated bands of a rotifer, pluteus and trochophore', *Biological Bulletin*, 142 (1972), 505-519.
- 181 Strathmann, note 171.
- 182 M.W. Hart, 'Deconstructing suspension feeders by analysis of film and video', *Invertebrate Biology*, 115 (1996), 185-190.
- 183 Hart, note 182.
- 184 Strathmann, note 171.
- 185 Strathmann, note 171.
- 186 Strathmann, note 172.
- 187 J.S. Ryland, 'Physiology and ecology of marine bryozoans', *Advances in Marine Biology*, 14 (1976), 285-443.
- 188 J.E. Winston, 'Feeding in marine bryozoans', in *Biology of Bryozoans*, edited by R.M. Woollacott and R.L. Zimmer (Academic Press, London, New York, 1977), 233-271.
- 189 Winston, note 3.
- 190 M. LaBarbera, 'Feeding currents and particle capture mechanisms in suspension feeding animals', *American Zoologist*, 24 (1984), 71-84.
- 191 F.K. McKinney and J.B.C. Jackson, *Bryozoan evolution* (University of Chicago Press, Chicago, London, 1989), 238.
- 192 M.W. Hart, 'Particle captures and the method of suspension feeding by echinoderm larvae', *Biological Bulletin*, 180 (1991), 12-27.
- 193 Hart, note 182.
- 194 Ryland, note 187.
- 195 Ryland, note 187, p. 327.
- 196 J.S. Ryland, 'Parameters of the lophophore in relation to population structure in a bryozoan community', in *Proceedings of the 9th European Marine Biology Symposium*, edited by H. Barnes (Aberdeen University Press, 1975), 363-393.
- 197 Ryland, note 187.
- 198 Ryland, note 196.
- 199 See also Cori, note 130.
- 200 For comparison see Markham and Ryland, note 133.
- 201 W.G. Sanderson, J.P. Thorpe and A. Clarke, 'Aspects of structure and function in the feeding of the Antarctic cheilostomate bryozoan, *Himantozoum antarcticum* (Calvet)', in *Proceedings of the 11th International Bryozoology Association Conference*, edited by A. Herrera Cubilla and J.B.C. Jackson (Smithsonian Tropical Research Institute, Panama, 2000), 355-364 (361).
- 202 Best and Thorpe, note 137.
- 203 Nielsen, note 178.
- 204 Sanderson, Thorpe and Clarke, note 201, p. 361.
- 205 See also M.A. Best and J.P. Thorpe, 'Feeding current interactions and competition for food among the bryozoan epiphytes of *Fucus serratus*', *Marine Biology*, 93 (1986), 371-375.
- 206 Best and Thorpe, note 137.
- 207 Borg, note 106.
- 208 Atkins, note 125.

- 209 Best and Thorpe, note 137, p. 91.
- 210 See for further discussion F.K. McKinney, 'Feeding and associated colonial morphology in marine bryozoans', *Critical Reviews in Aquatic Sciences*, 2(2) (1990), 255-280.
- 211 For comparison see Markham and Ryland, note 133.
- 212 Best and Thorpe, note 137.
- 213 Winston, note 188.
- 214 Winston, note 3.
- 215 J.E. Winston, 'Current-related morphology and behaviour in some Pacific coast bryozoans', in *Advances in Bryozoology*, edited by G.P. Larwood and M.B. Abbott, Systematics Association Special vol 13 (Academic Press, London, New York, 1979), 247-268.
- 216 Summarized in J.E. Winston, 'Feeding behaviour of modern bryozoans', in *Studies in Geology 5: Lophophorates. Notes for a Short Course*, edited by T.W. Broadhead (University of Tennessee, 1981), 1-21.
- 217 Winston, note 3.
- 218 Winston, note 188.
- 219 Bullivant, note 152.
- 220 Winston, note 3, p. 1.
- 221 Winston, note 3, p. 29.
- 222 M.H. Dick, 'Bryozoan behaviour in relation to autocleaning in *Holoporella brunea* (Hincks)', *Northwest Science*, 58(3) (1984), 195-207.
- 223 Winston, note 3, p. 13.
- 224 M.H. Dick, 'A proposed mechanism for chimney formation in encrusting bryozoan colonies', in *Bryozoa: Present and Past*, edited by J.R.P. Ross (Western Washington University, Bellingham, WA, 1987), 73-80 (74).
- 225 P.L. Cook, 'Bryozoa from Ghana', *Zoologische Wetenschappen Musee Royal l'Afrique centrale Tervuren, Belgique*, 238 (1985), 1-315.
- 226 Reviewed by McKinney, note 210.
- 227 Hunter and Hughes, note 138, p. 327, figure 2E.
- 228 Borg, note 106.
- 229 Strathmann, note 171.
- 230 Strathmann, note 172.
- 231 Hunter and Hughes, note 138.
- 232 H.U. Riisgård and P. Manríquez, 'Filter-feeding in fifteen marine ectoprocts (Bryozoa): particle capture and water pumping', *Marine Ecology Progress Series*, 154 (1997), 223-239.
- 233 C. Nielsen and H.U. Riisgård, 'Tentacle structure and filter-feeding in *Crisia eburnea* and other cyclostomatous bryozoans, with a review of upstream-collecting mechanisms', *Marine Ecology Progress Series*, 168 (1998), 163-186.
- 234 Rubenstein and Koehl, note 168.
- 235 LaBarbera, note 190.
- 236 J. Shimeta and P. Jumars, 'Physical mechanisms and rates of particle capture by suspension feeders', *Oceanography and Marine Biology Annual Review*, 29 (1991), 191-257.
- 237 Vogel, note 175.
- 238 Hart and Strathmann, note 176.
- 239 H.U. Riisgård and P.S. Larsen, 'Filter-feeding in marine macroinvertebrates: pump characteristics, modelling and energy cost', *Biological Review*, 70 (1996), 67-106.

- 240 H.U. Riisgård, P.S. Larsen and P.F. Nielsen, 'Particle capture in the mussel *Mytilus edulis*: the role of latero-frontal cirri', *Marine Biology*, 127 (1996), 259-266.
- 241 See for discussion also C.B. Jørgensen, 'A hydromechanical principle for particle retention in *Mytilus edulis* and other ciliary suspension feeders', *Marine Biology*, 61 (1981), 277-822.
- 242 C.B. Jørgensen, 'Fluid mechanics of the mussel gill: the lateral cilia', *Marine Biology*, 70 (1982), 275-281.
- 243 C.B. Jørgensen, 'Fluid mechanical aspects of suspension feeding', *Marine Ecology Progress Series*, 11 (1983), 89-103.
- 244 J. Shimeta and M.A.R. Koehl, 'Mechanisms of particle selection by tentaculate suspension feeders during encounter, retention, and handling', *Journal of Experimental Marine Biology and Ecology*, 209 (1997), 47-73.
- 245 D. Wildish and D.D. Kristmanson, *Benthic suspension feeders and flow* (Cambridge University Press, 1997), xiii, 409.
- 246 Riisgård and Manríquez, note 232, p. 237.
- 247 For further discussion see also Shunatova and Ostrovsky, note 1.
- 248 Nielsen, note 178.
- 249 Nielsen and Riisgård, note 233.
- 250 Riisgård and Manríquez, note 232.
- 251 Best and Thorpe, note 205.
- 252 W.G. Sanderson and J.P. Thorpe, 'Effects of temperature on the feeding activity of some temperate intertidal Bryozoa', in *Bryozoans in Space and Time*, edited by D.P. Gordon, A.M. Smith and J.A. Grant-Mackie (National Institute of Water and Atmospheric Research Ltd, Wellington, 1996), 271-281.
- 253 See similar opinion in B. Okamura, 'Particle size and flow velocity induce an inferred switch in bryozoan suspension-feeding behaviour', *Biological Bulletin*, 173 (1987), 222-229.
- 254 Shunatova and Ostrovsky, note 1.
- 255 Shunatova and Ostrovsky, note 1, p. 138.
- 256 I.I. Antipenko, *Feeding and behaviour of some fresh-water bryozoan species (Bryozoa, Phylactolaemata)*, Unpublished MSc Thesis, (St. Petersburg State University, 1998), 112 (in Russian).
- 257 Bullivant, note 162.
- 258 N.R. Menon, 'Clearance rates of food suspension and food passage rates as a function of temperature in two North-Sea bryozoans', *Marine Biology*, 24 (1974), 65-67.
- 259 J. Dudley, 'Differential utilization of phytoplankton food resources by marine entoprocts', *Biological Bulletin*, 139(2) (1970), 420.
- 260 Ryland, note 196.
- 261 See also Winston, note 188.
- 262 Winston, note 3.
- 263 Winston, note 215.
- 264 D. Jebram, 'Interrelations of nutrition, food uptake and growth in bryozoans', in *Advances in Bryozoology*, edited by G.P. Larwood and M.B. Abbott (Academic Press, London, 1979), 121-140.
- 265 J.P. Thorpe, D.R.K. Clarke and M.A. Best, 'Natural variation in tentacle number in marine bryozoans and the possible effects of intraspecific and interspecific ecological competition for food', in *Bryozoa: Ordovician to Recent*, edited by C. Nielsen and G.P. Larwood (Olsen

- and Olsen, Fredensborg, 1985), 319-327.
- 266 A. Herrera and J.B.C. Jackson, 'Environmental versus taxonomic variation in lophophore morphology of Panamanian cheilostome bryozoans', *Marine Biology*, 112 (1992), 99-106.
- 267 Best and Thorpe, note 137.
- 268 Best and Thorpe, note 205.
- 269 M.A. Best and J.P. Thorpe, 'Particle size, clearance rate and feeding efficiency in marine Bryozoa', in *Biology and Palaeobiology of Bryozoans*, edited by P.J. Hayward, J.S. Ryland and P.D. Taylor (Olsen and Olsen, Fredensborg, 1994a), 9-14.
- 270 W.G. Sanderson, J.P. Thorpe and A. Clarke, 'A preliminary study of feeding rates in the Antarctic cheilostomate bryozoan *Himantozoum antarcticum*', in *Biology and Palaeobiology of Bryozoans*, edited by P.J. Hayward, J.S. Ryland and P.D. Taylor (Olsen and Olsen, Fredensborg, 1994a), 167-172.
- 271 Sanderson and Thorpe, note 252.
- 272 Riisgård and Manríquez, note 232.
- 273 See also F.K. McKinney, M.R.A. Listokin and C.D. Phifer, 'Flow and polypide distribution in the cheilostome *Bugula* and their inference in *Archimedes*', *Lethaia*, 19 (1986), 81-93 (quoted in McKinney 1990).
- 274 Wildish and Kristmanson, note 245.
- 275 Reviewed in M.A. Best and J.P. Thorpe, 'The effect of suspended particulate matter (silt) on the feeding activity of the intertidal ctenostomate bryozoan *Flustrellidra hispida* (Fabricius)', in *Bryozoans in Space and Time*, edited by D.P. Gordon, A.M. Smith and J.A. Grant-Mackie (National Institute of Water and Atmospheric Research Ltd, Wellington, 1996), 39-45.
- 276 For discussion see also D.P. Gordon, A.G. Clark and J.F. Harper, 'Bryozoa', in *Animal Energetics*, edited by T.J. Pandian and F.J. Vernberg, volume 2 (London, Academic Press, 1987), 173-199.
- 277 Best and Thorpe, note 269.
- 278 Dudley, note 259.
- 279 Ryland, note 196.
- 280 J.S. Ryland, 'Physiology and ecology of marine bryozoans', *Advances in Marine Biology*, 14 (1976), 285-443.
- 281 Winston, note 188.
- 282 Winston, note 3, p. 1.
- 283 Winston, note 216.
- 284 Dudley, note 259.
- 285 Ryland, note 196.
- 286 Winston, note 188.
- 287 Winston, note 216.
- 288 McKinney and Jackson, note 191.
- 289 McKinney, note 210.
- 290 Herrera and Jackson, note 266.
- 291 Best and Thorpe, note 269.
- 292 Sanderson, Thorpe and Clarke, note 201, p. 361.
- 293 J.S. Porter, P. Bloor, B.A. Stokell and J.S. Ryland, 'Intra- and interspecific variation in tentacle number in the genus *Alcyonidium*', in *Proceedings of the 11th International Bryozoology Association Conference*, edited by A. Herrera Cubilla and J.B.C. Jackson

- (Smithsonian Tropical Research Institute, Panama, 2000), 321-328 (328).
- 294 Sanderson, Thorpe and Clarke, note 200, p. 361.
- 295 Reviewed in O.D. Hunt, 'The food of the bottom fauna of the Plymouth fishing grounds', *Journal of the Marine Biological Association of the United Kingdom*, New Series, 13 (1925), 560-599.
- 296 Jørgensen, note 151.
- 297 Bullivant, note 162.
- 298 Winston, note 188.
- 299 Winston, note 3.
- 300 Winston, note 215.
- 301 D. Jebram, 1977. 'Experimental techniques and culture methods', in *The Biology of Bryozoans*, edited by R.M. Woollacott and R.L. Zimmer (Academic Press, New York, London, 1977), 273-305.
- 302 McKinney and Jackson, note 191, p. 238.
- 303 B. Okamura, 'Particle size, flow velocity and suspension feeding by erect bryozoans *Bugula neritina* and *Bugula stolonifera*', *Marine Biology*, 105 (1990), 33-35.
- 304 M.A. Best and J.P. Thorpe, 'The uptake of ¹⁴C radioisotope tracer from dissolved glycine and glucose by the marine bryozoan *Flustrellidra hispida*', in *Bryozoaires Actuels et Fossiles: Bryozoa Living and Fossil*, edited by F.P. Bigey, *Bulletin de la Société des Sciences Naturelles de l'Ouest de la France*, Mémoire Hors Série, 1 (1991), 13-23.
- 305 M.A. Best and J.P. Thorpe, 'An analysis of potential food sources available to intertidal bryozoans in Britain', in *Biology and Palaeobiology of Bryozoans*, edited by P.J. Hayward, J.S. Ryland and P.D. Taylor (Olsen and Olsen, Fredensborg, 1994), 1-8.
- 306 Best and Thorpe, note 275.
- 307 Best and Thorpe, note 305.
- 308 D.K.A. Barnes and A. Clarke, 'Seasonal variation in the feeding activity of four species of Antarctic Bryozoa in relation to environmental factors', *Journal of Experimental Marine Biology and Ecology*, 181 (1994), 117-133.
- 309 D.K.A. Barnes and A. Clarke, 'Seasonality of feeding ecology in polar suspension feeders', *Polar Biology*, 15 (1995), 335-340.
- 310 W.G. Sanderson, S.P. Hardling and J.P. Thorpe, 'An investigation of the effects of tidal fluctuations on the availability of seston as food for intertidal Bryozoa', in *Bryozoans in Space and Time*, edited by D.P. Gordon, A.M. Smith and J.A. Grant-Mackie (Wellington, National Institute of Water and Atmospheric Research Ltd, 1996), 259-269.
- 311 B. Okamura, note 253.
- 312 Marcus, note 117.
- 313 N.R. Menon, 'Heat tolerance, growth and regeneration in three North Sea bryozoans exposed to different constant temperatures', *Marine Biology*, 15 (1972), 1-11.
- 314 N.R. Menon, note 258.
- 315 Riisgård and Manríquez, note 233.
- 316 Sanderson and Thorpe, note 252.
- 317 B. Okamura, 'The effects of ambient flow velocity, colony size, and upstream colonies on the feeding success of Bryozoa. I. *Bugula stolonifera* Ryland, an arborescent species', *Journal of Experimental Marine Biology and Ecology*, 83 (1984), 179-193.
- 318 B. Okamura, 'The effects of ambient flow velocity, colony size, and upstream colonies on the feeding success of Bryozoa. II. *Conopeum reticulum* (Linnaeus), an encrusting species',

- Journal of Experimental Marine Biology and Ecology*, 89 (1985), 69-80.
- 319 B. Okamura, 'Seasonal changes in zooid size and feeding activity in epifaunal colonies of *Electra pilosa*', in *Bryozoa: Present and Past*, edited by J.R.P. Ross (Western Washington University, Bellingham, WA, 1987), 197-203.
- 320 Okamura, note 303.
- 321 Okamura, note 303.
- 322 B. Okamura, 'The influence of neighbors on the feeding of an epifaunal bryozoan', *Journal of Experimental Marine Biology and Ecology*, 120 (1988), 105-123.
- 323 B. Okamura, 'Microhabitat variation and patterns of colony growth and feeding in a marine bryozoan', *Ecology*, 73 (1992), 1502-1513.
- 324 L.W. Buss, 'Bryozoan overgrowth interactions – the interdependence of competition for space and food', *Nature*, 281(5731), (1980), 475-477.
- 325 Best and Thorpe, note 205.
- 326 M.A. Best and J.P. Thorpe, 'Effects of food particle concentration on feeding current velocity in six species of marine Bryozoa', *Marine Biology*, 93 (1986), 255-262.
- 327 See also L.W. Buss, 'Mechanisms of competition between *Onychozella* (Hastings) and *Antropora tincta* (Hastings) on an Eastern Pacific rocky shoreline', in *Recent and Fossil Bryozoa*, edited by G.P. Larwood and C. Nielsen (Olsen and Olsen, Fredensborg, 1981) 39-49.
- 328 L.W. Buss and J.B.C. Jackson, 'Planctonic food availability and suspended feeder abundance; evidence of in situ depletion', *Journal of Experimental Marine Biology and Ecology*, 49 (1981), 151-161.
- 329 Thorpe, Clarke and Best, note 265.
- 330 P.S. Larsen, S.S. Matlok and H.U. Riisgård, 'Bryozoan filter feeding in laminar wall layers: flume experiments and computer simulation', *Vie Milieu*, 48(4) (1998), 309-319.
- 331 J.E. Eckman and B. Okamura, 'A model of particle capture by bryozoans in turbulent flow: significance of colony form', *American Naturalist*, 152(6) (1998), 861-880.
- 332 J.W. Bishop and L.M. Bahr, 'Effects of colony size on feeding by *Lophopodella carteri* (Hyatt)', in *Animal Colonies: Development and Function through Time*, edited by R.S. Boardman, A.H. Cheetham and W.A. Oliver (Dowden, Hutchinson and Ross, Stroudsburg, PA, 1973), 433-437.
- 333 B. Okamura and L.A. Doolan, 'Patterns of suspension feeding in the freshwater bryozoan *Plumatella repens*', *Biological Bulletin*, 184 (1993), 52-56.
- 334 Winston, note 3, p. 1.
- 335 Okamura, note 319.
- 336 McKinney, note 210.
- 337 Riisgård and Manríquez, note 232.
- 338 Porter, Bloor, Stokell and Ryland, note 293.
- 339 Sanderson, Thorpe and Clarke, note 201, p. 361.
- 340 Shunatova and Ostrovsky, note 1.
- 341 See as examples Riisgård and Manríquez, note 232.
- 342 Nielsen and Riisgård, note 233.
- 343 Sanderson, Thorpe and Clarke, note 201, p. 361.

